

Name :

Roll No. :

Invigilator's Signature :

CS/B.Tech(N)/SEM-1/M-101/2012-13

2012

MATHEMATICS-I

Time Allotted : 3 Hours

Full Marks : 70

The figures in the margin indicate full marks.

*Candidates are required to give their answers in their own words
as far as practicable*

GROUP - A

(Multiple Choice Type Questions)

1. Choose the correct alternatives for any *ten* of the following :

10 × 1 = 10

i) The sequence $\left\{ (-1)^n \frac{1}{n} \right\}$ is

a) Convergent

b) Oscillatory

c) Divergent

d) none of these.

ii) The matrix $\begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$ is

a) Symmetric

b) Skew-symmetric

c) Singular

d) Orthogonal.

1151 (N)

[Turn over

CS/B.Tech(N)/SEM-1/M-101/2012-13

GROUP - C**(Long Answer Type Questions)**Answer any *three* of the following. $3 \times 15 = 45$

7. a) If $u = x^2 - 2y$, $v = x + y + z$, $w = x - 2y + 3z$,
find $\frac{\partial (u, v, w)}{\partial (x, y, z)}$.

b) Prove that
$$\begin{vmatrix} 1 & \alpha & \alpha^2 - \beta\gamma \\ 1 & \beta & \beta^2 - \gamma\alpha \\ 1 & \gamma & \gamma^2 - \alpha\beta \end{vmatrix} = 0.$$

- c) If $v = f(x^2 + 2yz, y^2 + 2zx)$, prove that

$$(y^2 - zx) \frac{\partial v}{\partial x} + (x^2 - yz) \frac{\partial v}{\partial y} + (z^2 - xy) \frac{\partial v}{\partial z} = 0.$$

5 + 5 + 5

8. a) If $\theta = t^n e^{-\frac{r^2}{4t}}$, find what value of n will make
 $\frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial \theta}{\partial r} \right) = \frac{\partial \theta}{\partial t}$.

- b) Using mean value theorem prove that

$$0 < \frac{1}{x} \log \left(\frac{e^x - 1}{x} \right) < x.$$

- c) If $I_n = \int_0^{\frac{\pi}{2}} x^n \sin x \, dx$ ($n > 1$), then show that

$$I_n + n(n-1)I_{n-2} = n \left(\frac{\pi}{2} \right)^{n-1}. \quad 5 + 5 + 5$$

CS/B.Tech(N)/SEM-1/M-101/2012-13

9. a) State D'Alembert's ratio test for convergence of an infinite series. Examine the convergence or divergence of the series $\left(\frac{1}{3}\right)^2 + \left(\frac{1.2}{3.5}\right)^2 + \left(\frac{1.2.3}{3.5.7}\right)^2 + \dots$

b) If $y = e^{\tan^{-1}x}$, then show that $(1 + x^2) y_{n+2} + (2nx + 2x - 1) y_{n+1} + n(n+1) y_n = 0$.

c) Find the extreme value of the function

$$f(x, y) = x^3 + y^3 - 3x - 12y + 20. \quad 5 + 5 + 5$$

10. a) If $A = \begin{bmatrix} 1 & 0 & 2 \\ 0 & -1 & 1 \\ 0 & 1 & 0 \end{bmatrix}$, then verify that A satisfies its

own characteristic equation. Hence find A^{-1} and A^9 .

b) If $u = \tan^{-1} \left(\frac{x^3 + y^3}{x - y} \right)$, then show that

$$x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} = (1 - 4 \sin^2 u) \sin 2u.$$

c) Given the system of equation :

$$x_1 + 4x_2 + 2x_3 = 1, \quad 2x_1 + 7x_2 + 5x_3 = k,$$

$$4x_1 + mx_2 + 10x_3 = 2k + 1. \text{ Find for what values of}$$

k and m , the system has (i) an unique solution,

(ii) no solution (iii) many solution.

CS/B.Tech(N)/SEM-1/M-101/2012-13

11. a) Show that $\vec{\nabla} r^n = nr^{n-2} \vec{r}$,

$$\text{where } \vec{r} = \vec{i}x + \vec{j}y + \vec{k}z.$$

b) Evaluate $\int \int \sqrt{4x^2 - y^2} \, dx dy$ over the triangle formed by the straight lines $y = 0$, $x = 1$ and $y = x$.

c) Verify Stokes theorem for

$\vec{F} = (2x - y) \hat{i} - yz^2 \hat{j} - y^2 z \hat{k}$, where S is the upper half surface of the sphere $x^2 + y^2 + z^2 = 1$ and C is its boundary. 5 + 5 + 5
